



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

173. Proposed by P. C. CULLEN, Principal of Schools, Indianola, Neb.

To construct a circle tangent to a given line at a given point such that tangents drawn to this circle and passing through two fixed points shall be parallel.


No solution of this problem has been received.

174. Proposed by J.M.HOWIE, Professor of Mathematics, The Nebraska State Normal School, Peru, Neb.

Describe a circle which shall pass through a given point and be tangent to two given circles.

Solution by G. B. M. ZERR, A. M., Ph. D., The Temple College, Philadelphia, Pa., and M. E. GRABER, Heidelberg University, Tiffin, Ohio.

Let C , C' be the centers of the given circles; E the given point. Describe a circle through EBB' and let H be the homothetic center of C , C' . Draw FEH and $B'BH$. Let the circle through $B'BE$ cut C again in R . Draw BRP and from P draw PD tangent to C , then the circle circumscribing the triangle FED is the circle required, for $HD \cdot HG = HB' \cdot HB$. Since there can be two tangents drawn from P to C , there can be drawn two circles tangent to C , C' so that the line joining their points of contact shall pass through H . Similarly, two circles can be drawn tangent to C , C' passing through E so that the line joining their points of contact shall pass through the anti-homothetic center H' lying between C and C' .



Therefore, there are four circles satisfying the condition.

175. Proposed by W. P. WEBBER, Mississippi Normal College, Houston, Miss.

A field is enclosed by a fence in circular form and a straight gate 20 feet wide. The fence is 100 feet in length. How much land is in the field? [Solution by most elementary method possible.]

Solution by G. B. M. ZERR, A. M., Ph. D., The Temple College, Philadelphia, Pa.; J. SCHEFFER, A. M., Hagerstown, Md.; and the PROPOSER.

Chord $AB = 20$, arc $ADB = 100^\circ$.

Let $\angle ACB = \theta$, $AC = r$.

Then $\lceil 2\pi - \theta \rceil r = 100$, $r \sin \frac{1}{2}\theta = 10$.

$$\therefore 2\pi = \theta + 10 \sin \frac{1}{2}\theta.$$

$$\therefore \theta = 62^\circ 32' \text{ nearly, } r = 19.267 \text{ feet.}$$

$$\text{Area sector } ADB = \frac{[360^\circ - 62^\circ 32']\pi r^2}{360^\circ}$$

\equiv 963.64 square feet.

Area triangle $ACB = 164.69$ square feet.

Total area = 1128.33 square feet.

